

SOLAR-LIKE DIFFERENTIAL ROTATION IN A CONVECTIVE DYNAMO WITH A CORONAL ENVELOPE

JÖRN WARNECKE^{1,2}, PETRI J. KÄPYLÄ^{1,3}, MAARIT J. MANTERE^{3,4} AND AXEL BRANDENBURG^{1,2}¹NORDITA, KTH Royal Institute of Technology and Stockholm University, Roslagstullsbacken 23, SE-10691 Stockholm, Sweden²Department of Astronomy, AlbaNova University Center, Stockholm University, SE-10691 Stockholm, Sweden³Physics Department, Gustaf Hållströmin katu 2a, PO Box 64, FI-00014 University of Helsinki, Finland⁴Aalto University, Department of Information and Computer Science, PO Box 15400, FI-00076 Aalto, Finland*Draft version January 11, 2013*

ABSTRACT

We report on the results of three convective dynamo simulations with an outer coronal layer. The magnetic field is self-consistently generated by the convective motions beneath the surface. Above the convection zone we include a polytropic layer that extends to 1.6 solar radii. The temperature increases in this region to ≈ 8 times the value at the surface, corresponding to ≈ 1.2 times the value at the bottom of the spherical shell. We associate this region with the solar corona. We find a solar-like differential rotation with radial contours of constant rotation rate, together with a solar-like meridional circulation and a near-surface shear layer. This spoke-like rotation profile is caused by a non-zero latitudinal entropy gradient which violates the Taylor-Proudman balance via the baroclinic term. The lower density stratification compared with the Sun leads to an equatorward return flow above the surface. The mean magnetic field is in most of the cases oscillatory with equatorward migration in one case. In other cases the equatorward migration is overlaid by stationary or even poleward migrating mean fields.

Subject headings: Magnetohydrodynamics – convection – turbulence – Sun: dynamo – Sun: rotation – Sun: activity

1. INTRODUCTION

The Sun has an activity cycle of eleven years, which is manifested by sunspots occurring at the solar surface. The sunspot number changes from a few during minimum to over 200 during maximum. The sunspot locations display a latitudinal dependence during the cycle. At solar minimum, sunspots emerge preferably at higher latitudes and during maximum at lower latitudes. By plotting these for several cycles, one obtains the ‘butterfly diagram’. Every eleven years the polarity of the sunspot pairs changes sign, manifesting the 22-year magnetic cycle. To understand this cyclic behavior, one has to connect the fluid motions in the Sun with magnetic field generation to construct dynamo models. These dynamo models should be able to reproduce the 22-year magnetic activity cycle as well as the large-scale magnetic field evolution at the surface of the Sun. It is widely believed that the sunspots are correlated with the large-scale magnetic field distribution. Therefore, a successful solar dynamo model should reproduce the equatorward migration of the large-scale field as we observe it indirectly from sunspots and more directly from synoptic magnetograms.

Until recently, only kinematic mean-field models, where turbulent effects are parameterized through transport coefficients (e.g. Krause & Rädler 1980), have been able to show this (e.g. Dikpati & Charbonneau 1999; Käpylä et al. 2006; Kitchatinov & Olemskoy 2012). These models have been used to reproduce certain features of the solar cycle, such as the Maunder minimum (e.g., Karak 2010). However, these models are only valid in the kinematic regime where the fluid motions are assumed given, so they are not self-consistently generated. The backreaction from the magnetic field is either ignored or taken into account in a rudimentary way involving ad hoc quenching for the turbulent trans-

port coefficients. Direct numerical simulations (DNS) of the solar dynamo have been unsuccessful in producing equatorward migration using convective motions to drive a dynamo (e.g. Gilman 1983; Brun et al. 2004; Käpylä et al. 2010; Ghizaru et al. 2010; Brown et al. 2011) so far. This was possibly due to the low fluid and magnetic Reynolds numbers in the simulations. Equatorward migration of the mean magnetic field was, for the first time, found in DNS by Käpylä et al. (2012). The exact cause is not yet fully understood, but the amount of density stratification seems to play an important role (Käpylä et al. 2013).

An important ingredient of the solar dynamo is differential rotation. It is believed that strong shear at the bottom of the convection zone (Spiegel & Weiss 1980) or near the surface (Brandenburg 2005), plays an important role amplifying the magnetic field. However, even today it is not straightforward to reproduce a solar-like differential rotation profile. Mean-field simulations (Brandenburg et al. 1992; Kitchatinov & Rüdiger 1995) have been able to reproduce a solar-like rotation profile by modeling small-scale effects by mean-field coefficients through the Λ effect (see e.g. Rüdiger 1980, 1989). These models reproduce the positive (negative) latitudinal gradient of angular velocity in the northern (southern) hemisphere – i.e., the equator rotates faster than the poles – together with ‘spoke-like’ contours in the meridional plane. DNS of convective dynamos are able to reproduce a rapidly rotating equator at sufficiently large Coriolis numbers (Brun et al. 2004; Käpylä et al. 2011b). To generate spoke-like differential rotation in DNS, however, has so far been reported to be possible only by imposing a latitudinal entropy gradient at the bottom of the convection zone (Miesch et al. 2006). A self-consistently generated spoke-like profile has not yet been found.

An important issue with solar dynamo models is the effect of catastrophic quenching of the dynamo at high mag-

netic Reynolds numbers; see Brandenburg & Subramanian (2005). This is caused by the accumulation of magnetic helicity in the dynamo region. DNS provide evidence that magnetic helicity fluxes both within and through the boundaries of the dynamo domain can prevent the dynamo from being catastrophically quenched (e.g. Brandenburg & Sandin 2004; Hubbard & Brandenburg 2012). In the case of the Sun, magnetic helicity flux can emerge through the solar surface and can be transported away from the Sun by coronal mass ejections or by the solar wind (Blackman & Brandenburg 2003). In earlier work we mimic the case of the Sun by using an upper layer on the top of a dynamo region to allow for magnetic helicity fluxes leaving the domain (Warnecke & Brandenburg 2010; Warnecke et al. 2011, 2012a,b). This two-layer model was successful in showing that the dynamo is not only enhanced, but it can actually trigger the emergence of coronal ejections. These ejections have a similar shape as coronal mass ejections and carry a significant amount of magnetic helicity out of the dynamo region.

In this work we use the two-layer approach to investigate the influence of the coronal layer as an upper boundary condition for a convective dynamo. We focus on the physical properties and dynamics in the convection zone. The effects of varying the strength of stratification on a convective dynamo without corona is studied in a companion paper (Käpylä et al. 2013).

2. MODEL AND SETUP

We use a two-layer model in spherical polar coordinates (r, θ, ϕ) , where the lower layer ($r \leq R$) represents the convection zone and the upper one the corona. The simulation is performed in a spherical wedge with radial extent $r_0 \leq r \leq R_c = 1.6R$, where $r_0 = 0.7R$ corresponds to the bottom of the convection zone and R to the solar radius, for colatitudes $15^\circ \leq \theta \leq 165^\circ$ and an azimuthal extent $0 \leq \phi \leq 45^\circ$. We solve the following equations of compressible magnetohydrodynamics,

$$\frac{\partial \mathbf{A}}{\partial t} = \mathbf{u} \times \mathbf{B} + \eta \nabla^2 \mathbf{A}, \quad (1)$$

$$\frac{D \ln \rho}{Dt} = -\nabla \cdot \mathbf{u}, \quad (2)$$

$$\frac{D\mathbf{u}}{Dt} = \mathbf{g} - 2\boldsymbol{\Omega}_0 \times \mathbf{u} + \frac{1}{\rho} (\mathbf{J} \times \mathbf{B} - \nabla p + \nabla \cdot 2\nu\rho\mathbf{S}), \quad (3)$$

$$T \frac{Ds}{Dt} = \frac{1}{\rho} \nabla \cdot (\mathbf{F}^{\text{rad}} + \mathbf{F}^{\text{SGS}}) + 2\nu\mathbf{S}^2 + \frac{\mu_0\eta}{\rho} \mathbf{J}^2 - \Gamma_{\text{cool}}(r), \quad (4)$$

where the magnetic field is given by $\mathbf{B} = \nabla \times \mathbf{A}$ and thus obeys $\nabla \cdot \mathbf{B} = 0$ at all times, $\mathbf{J} = \mu_0^{-1} \nabla \times \mathbf{B}$ is the current density, μ_0 is the vacuum permeability, η and ν are the magnetic diffusivity and kinematic viscosity, respectively, $D/Dt = \partial/\partial t + \mathbf{u} \cdot \nabla$ is the advective time derivative, ρ is the density, and $\mathbf{u}$ is the velocity. The traceless rate-of-strain tensor is given by

$$S_{ij} = \frac{1}{2}(u_{i;j} + u_{j;i}) - \frac{1}{3}\delta_{ij} \nabla \cdot \mathbf{u}, \quad (5)$$

where semicolons denote covariant differentiation; see Mitra et al. (2009) for details. Furthermore, $\boldsymbol{\Omega}_0 = \Omega_0(\cos\theta, -\sin\theta, 0)$ is the rotation vector and p is the pressure. The gravitational acceleration is given by

$$\mathbf{g} = -GM\mathbf{r}/r^3, \quad (6)$$

where G is Newton's gravitational constant and M is the mass of the star. The radiative and subgrid scale (SGS) heat fluxes are defined as

$$\mathbf{F}^{\text{rad}} = K\nabla T, \quad \mathbf{F}^{\text{SGS}} = \chi_{\text{SGS}}\rho T\nabla s, \quad (7)$$

where K is the radiative heat conductivity and χ_{SGS} turbulent heat conductivity, which represents the unresolved convective transport of heat. The fluid obeys the ideal gas law, $p = (\gamma - 1)\rho e$, where $\gamma = c_P/c_V = 5/3$ is the ratio of specific heats at constant pressure and constant volume, respectively, and $e = c_V T$ is the internal energy density, defining the temperature T . Finally, $\Gamma_{\text{cool}}(r)$ is the radial cooling profile which is specified in Equation (10).

The two-layer model is similar to that used in the previous work (Warnecke & Brandenburg 2010; Warnecke et al. 2011, 2012a,b), except that here we improve the model of Warnecke et al. (2012b) in two important ways. First, we use a more realistic model for the convection zone as in Käpylä et al. (2011a, 2012). Instead of using a polytropic setup with $m = 1$, we lower the radiative flux by decreasing K and introducing a turbulent heat conductivity χ_{SGS} , (referred to as χ_t in Käpylä et al. 2012). We apply a piecewise constant profile for χ_{SGS} such that in the interval of $0.75R \leq r \leq 0.97R$ it is equal to a quantity $\bar{\chi}_{\text{SGS}}$ (whose value is similar to that of the microphysical viscosity ν), and it goes smoothly to zero above and below the boundaries of the interval. Additionally, we change the temperature profile compared to our earlier isothermal cold corona to a temperature-stratified corona, which is ≈ 8 times hotter than the surface and ≈ 1.2 times than bottom of the convection zone. In Figure 1, the averaged temperature profile is shown together with the averaged density, pressure and entropy profiles for one of the runs.

We initialize the simulations with pre-calculated radial profiles of temperature, density, and pressure. In the convection zone ($r \leq R$) we have an isentropic and hydrostatic initial state for the temperature, whose gradient is given by

$$\frac{\partial T}{\partial r} = \frac{-|g(r)|}{c_V(\gamma - 1)(m + 1)}, \quad (8)$$

where $m = 1.5$ is the polytropic index. This leads to a temperature minimum T_{min} above the surface of the convective layer at $r = R$. In the corona ($R \leq r \leq R_c$), we prescribe the temperature as

$$T(r) = T_{\text{min}} + \frac{1}{2} (T_{\text{cor}} - T_{\text{min}}) \left[1 + \tanh \left(\frac{r - r_{\text{tra}}}{w} \right) \right], \quad (9)$$

where T_{cor} is temperature in the corona, and r_{tra} and the width $w = 0.02R$ are chosen to have a smooth temperature profile as shown in Figure 1. The radial cooling profile $\Gamma_{\text{cool}}(r)$ in Equation (4) maintains the temperature profile,

$$\Gamma_{\text{cool}} = \Gamma_0 f(r) (c_s^2 - c_{s0}^2)/c_{s0}^2, \quad (10)$$

where $f(r)$ is a profile function equal to unity in $r > R$ and going smoothly to zero in $r \leq R$, and Γ_0 is a cooling luminosity chosen so that the sound speed in the corona relaxes towards the temperature profile given in Equation (9). The stratification of density follows from the hydrostatic equilibrium. The radial heat conductivity profile is chosen so that the energy in the convection zone is mostly transported by convective motion. We apply a profile for the viscosity ν




FIG. 1.— Radial profiles of stratification for Run A. The normalized density ρ/ρ_0 (dashed lines), pressure p/p_0 (dash-triple-dotted), temperature T/T_0 (solid) are plotted together with the specific entropy s/c_p (dash-dotted) over the radius. The inset shows various profiles in logarithmic representation to emphasize the steep decrease of the pressure and density in the coronal layer. The index 0 represents the value at the bottom of the convection zone.

which is constant in the convection zone ($r \leq R$) and increase smoothly above the surface to a value that is 20 times higher in the corona. This helps to suppress high velocities and sharp flow structures aligned with the rotation vector in the corona especially in the beginning of the simulation, when the magnetic field is weak. We initialize the magnetic field with weak Gaussian-distributed perturbations inside the convection zone.

We use periodic boundary conditions in the azimuthal direction. For the velocity field we apply stress-free boundary condition at the radial and latitudinal boundaries. The magnetic field follows a perfect conductor condition at the lower radial and the two latitudinal boundaries. On the outer radial boundary, we force the field to be radial; for discussion on the applicability of this boundary condition for the Sun see Warnecke et al. (2012b). We set the gradient of the temperature at the lower radial boundary, corresponding to a fixed radiative flux, and at the radial outer boundary we set the temperature to a constant value. At the latitudinal boundaries, we impose a vanishing first derivative of entropy to suppress a heat flux through the boundary.

Our runs are characterized by the values of the fluid and magnetic Reynolds numbers, $Re = u_{rms}/\nu k_f$ and $Rm = u_{rms}/\eta k_f$, respectively, where u_{rms} is the volume averaged rms velocity in the convection zone, and $k_f = 2\pi/(R - r_0) \approx 21/R$ is used as a reference wavenumber. To represent the turbulent velocities in a proper way, we define $u_{rms} = \sqrt{3/2 \langle u_r^2 + u_\theta^2 \rangle_{\theta\phi r \leq R}}$ where we neglect the differential rotation dominated ϕ -component. We also define the fluid and magnetic Prandtl numbers $Pr = \nu/\chi_{SGS}$ and $Pm = \nu/\eta = Rm/Re$, the Coriolis number $Co = 2\Omega_0/u_{rms}k_f$ and Taylor number $Ta = (2\Omega_0 R^2/\nu)^2$. Time is given in turnover times, $\tau = (u_{rms}k_f)^{-1}$. We measure the magnetic field strength as the rms value averaged over the convection zone B_{rms} , and we normalize this value with the equipartition value of the magnetic field defined by $B_{eq}^2 = \mu_0 \langle \rho u^2 \rangle_{r \leq R}$. We use the (semi-) turbulent Rayleigh number Ra_t from the

thermally relaxed state of the run,

$$Ra_t = \frac{GM(R - r_0)^4}{\nu \chi_{SGS} R^2} \left(-\frac{1}{c_p} \frac{ds}{dr} \right)_{r=0.85 R}. \quad (11)$$

To monitor the solutions in the convection zone, we use two different heights, one near the surface at $r_1 = 0.97 R$, and one in the middle of the convection zone at $r_2 = 0.84 R$. We use the PENCIL CODE¹ with sixth-order centered finite differences in space and a third-order Runge-Kutta scheme in time; see Mitra et al. (2009) for the extension of the PENCIL CODE to spherical coordinates.

3. RESULTS

We focus in this work on three simulations which are summarized in Table 1. The main differences between these runs is their rotation rate and the magnetic Reynolds number. Run Ac is a continuation of Run A after $t/\tau = 1150$, but with a smaller diffusivity η in the convection zone. The Runs A and Ac have a higher Coriolis number Co and lower fluid and magnetic Reynolds number Re and Rm than Run B. We show the time evolution of the total rms velocity and magnetic field, averaged over the whole domain, $u_{rms}^{tot} = \langle u_r^2 + u_\theta^2 + u_\phi^2 \rangle_{r\theta\phi}^{1/2}$ and $B_{rms}^{tot} = \langle B_r^2 + B_\theta^2 + B_\phi^2 \rangle_{r\theta\phi}^{1/2}$, in all the three runs in Figure 2. Here, the subscripts on angle brackets denote averaging over r, θ, ϕ . Convection is sufficiently supercritical to develop during the first few tens of turnover times. After 50–200 turnover times, the dynamo starts to operate and a magnetic field grows at a rate that is higher for faster rotation (Run A). Due to high rotation rate and the lower density in the corona, the velocities there grow to higher values than in the convection zone. As we describe in Section 2, we use a higher viscosity to suppress these velocities and associated numerical difficulties. After the magnetic field in the convection zone has reached sufficient strength and expanded throughout the whole domain, it quenches the high velocities in the corona significantly, as evident from Figure 2. When the magnetic field reaches $B_{rms}^{tot} \approx 0.3 B_{eq}$ the rms velocity decreases from $u_{rms}^{tot} \approx 6$ to ≈ 1 , i.e., the contribution from the corona is now sub-dominant. This is caused by the Lorentz force, which becomes much stronger and comparable to the Coriolis force in the corona when the magnetic field has expanded into the whole domain. In the saturated state $u_{rms}^{tot} \approx u_{rms}$, which is reached after around $t/\tau = 1000$ turnover times for Runs A and Ac, even though the magnetic field is still growing slowly. For Run B, at first it seems that the saturated state has been reached at $t/\tau = 1000$, but it turns out that both u_{rms}^{tot} and magnetic field start growing again. While the differential rotation profile remains roughly unchanged despite of the growth of the energies (see Section 3.1), the magnetic field seems to undergo a mode change from oscillatory into a stationary solution or an oscillatory solution with a much longer period (see Section 3.2).

In Figure 3 we show the balance of various energy fluxes, contributing to the total luminosity of the simulated star for Run A. The radial components of radiative, convective, kinetic, viscous, and Poynting fluxes, as well as the flux due to the turbulent heat conductivity, are defined as

$$\overline{F}_{rad} = -\langle F_r^{rad} \rangle, \quad (12)$$

$$\overline{F}_{conv} = c_p \langle (\rho u_r)' T' \rangle, \quad (13)$$

¹ <http://pencil-code.googlecode.com>

TABLE 1
SUMMARY OF THE RUNS.

Run	grid	Pr	Pm	Ta	ρ_0/ρ_s	Ma	Ra_t	Re	Rm	Co	B_{rms}^2/B_{eq}^2	Δ_Ω
A	$400 \times 256 \times 192$	5	1	$1.4 \cdot 10^{10}$	14	0.08	$2.6 \cdot 10^6$	25	25	11	0.29	-0.01
Ac	$400 \times 256 \times 192$	5	1.67	$1.4 \cdot 10^{10}$	14	0.08	$2.6 \cdot 10^6$	25	42	11	0.27	-0.03
B	$400 \times 256 \times 192$	4	1	$7.2 \cdot 10^9$	15	0.09	$2.6 \cdot 10^6$	36	36	5.3	0.37	-0.05

NOTE. — The second to fourth columns show quantities that are input parameters to the models whereas the quantities in the last four columns are results of the simulations computed from the saturated state. The Mach number is defined as $Ma = u_{rms}/c_s|_{r=0.97R}$ and latitudinal differential rotation is quantified through $\Delta_\Omega = \partial\bar{\Omega}/\partial\cos^2\theta|_{r \leq R}/\Omega_0$.




FIG. 2.— Time evolution of the total rms velocity and magnetic field. The rms velocity of the whole domain u_{rms}^{tot} is normalized by u_{rms} (solid line) and is plotted together with the rms magnetic field of the whole domain B_{rms}^{tot} normalized by the equipartition field in the convection zone, B_{eq} , (dotted line) and multiplied by ten for visualization purposes for Run A (black line), Run Ac (blue), and Run B (red).

$$\overline{F}_{kin} = -\frac{1}{2} \langle (\rho u_r)' u'^2 \rangle, \quad (14)$$

$$\overline{F}_{visc} = -2\nu \langle (\rho u_i)' S_{ir}' \rangle, \quad (15)$$

$$\overline{F}_{SGS} = -\langle F_r^{SGS} \rangle, \quad (16)$$

$$\overline{F}_{Poynt} = \langle E_\theta B_\phi - E_\phi B_\theta \rangle / \mu_0, \quad (17)$$

where $\mathbf{E} = \eta\mu_0\mathbf{J} - \mathbf{u} \times \mathbf{B}$ is the electric field, the primes denote fluctuations, and angle brackets averaging over θ and ϕ and a time interval over which the turbulence is statistically stationary. The convective flux dominates inside the convection zone and reaches here much higher values than in our earlier model (Warnecke et al. 2012b). (In our case, $\overline{\chi}_{SGS} \approx 0.02\chi_{t0}$, where $\chi_{t0} = u_{rms}/3k_f$ is an estimate for the macrophysical turbulent diffusivity.) In the corona the cooling flux keeps the total flux constant. Note the convective overshoot into the exterior and the negative radiative flux just above the surface ($r = R$), caused by the higher temperature in the corona. The kinetic energy flux has small negatives values in the convection zone. The luminosity due to viscosity as well as the Poynting flux is too small to be visible.

3.1. Differential rotation and meridional circulation

In Figure 4, we plot the mean rotation profiles $\bar{\Omega}(r, \theta) = \Omega_0 + \bar{u}_\phi/r \sin\theta$ together with the meridional circulation $(\bar{u}_r, \bar{u}_\theta, 0)$ of Runs A, Ac and B. The contours of constant rotation are clearly not cylindrical. They show a ‘spoke-like’ structure, i.e. the contours are more radial than cylindrical, which is similar to the solar rotation profile obtained by helioseismology. The equator is rotating faster than the poles, which has been seen in previous simulations of Brun et al.




FIG. 3.— The different contributions to the total luminosity (solid purple line) are due to radiative diffusion (dashed red line), resolved convection (blue dotted), unresolved turbulent convection (black dotted), viscosity (yellow dashed), flux correspond to the cooling function Γ_{cool} (dash-dotted), and the Poynting flux (orange dash-dotted) for Run A. The thin solid black lines denote the zero level and the total luminosity through the lower boundary, respectively.

(2004), Miesch et al. (2006), and Käpylä et al. (2010, 2011b) and resembles the observed rotation of the Sun for our slower rotating case (Run B).

To produce spoke-like rotation contours, the Taylor–Proudman theorem has to be overcome by an important contribution in the evolution equation for the mean azimuthal vorticity $\overline{\omega}_\phi$, which is given by:

$$\frac{\partial \overline{\omega}_\phi}{\partial t} = r \sin\theta \frac{\partial \overline{\Omega}^2}{\partial z} + (\nabla T \times \nabla s)_\phi + \dots \quad (18)$$

where $\partial/\partial z = \cos\theta \partial/\partial r - r^{-1} \sin\theta \partial/\partial\theta$ is the derivative along the rotation axis. The first term in Equation (18) is related to the curl of the Coriolis force and vanishes for cylindrical $\overline{\Omega}$ contours. The second term is the mean baroclinic term, which is caused mainly by latitudinal entropy variations. We ignore here additional contributions such as meridional Reynolds and Maxwell stresses which turn out to be small. In Figure 5, we plot the first and second terms of Equation (18) for each of the three runs. These two contributions balance each other nearly perfectly. This lets us conclude that these two terms give the dominant contribution to the production of mean azimuthal vorticity, and that the Taylor–Proudman theorem is broken by the baroclinic term. It is remarkable that there is such a large and spatially coherent latitudinal entropy gradient, which is crucial to having a significant azimuthal baroclinic term, is produced self-consistently in the simulations.

Similar results have so far only been obtained in mean-field




FIG. 4.— Differential rotation and meridional circulation zoomed in the northern hemisphere in the convection zone for the three different Runs A, Ac, and B, from the top row to the bottom one. *Left column*: mean rotation profiles $\bar{\Omega}(r, \theta)/\Omega_0$ color coded and as rotation contours. *Middle column*: mean rotation profiles (color coded) overplotted with vectors (white) showing the meridional circulation in terms of the mass flux $\bar{\rho}(\bar{u}_r, \bar{u}_\theta, 0)$. Note the different plotting region in the first and middle columns. The white dotted lines indicate the radii $r = r_1 = 0.97 R$ and $r = r_2 = 0.84 R$. The three black dash-dotted lines represent the latitudes $90^\circ - \theta_1 = 30^\circ$, $90^\circ - \theta_2 = 15^\circ$, and $90^\circ - \theta_3 = 7^\circ$, which are used in the right column. The white-black dashed line indicates the surface ($r = R$). *Right column*: latitudinal mass flux $\bar{\rho} u_\theta/\rho_0 u_{rms}$ plotted over radius r/R for three different latitudes θ_1 (blue dashed line), θ_2 (red solid line) and θ_3 (black solid line). ρ_0 represents the value at the bottom of the convection zone. The dotted and dash-triple-dotted lines represent the corresponding values in the southern hemisphere; see the legend for details. The black dashed lines indicate the zero line, the surface ($r = R$) and the depths, r_1 and r_2 .

simulations (Brandenburg et al. 1992; Kitchatinov & Rüdiger 1995) by including an anisotropic convective heat conductivity, and in convection simulations by prescribing a latitudinal entropy gradient at the lower radial boundary of the convection zone (Miesch et al. 2006). The last column of Figure 5

shows the mean latitudinal entropy gradient $\nabla_\theta \bar{s}$ for the three runs. The spatial distribution of the gradient agrees with the baroclinic term as well, because $\nabla_r \bar{T} \approx \text{const}$, so we can conclude that the dominant contribution in the baroclinic term is due to the product of the latitudinal entropy gradient and the




FIG. 5.— Representation of the two dominant terms in the balance of mean azimuthal vorticity together with the latitudinal entropy gradient. In the first column the mean azimuthal component of the baroclinic term of the total entropy gradient and total temperature gradient $(\nabla T \times \nabla s)_\phi$ is plotted together with advection term $r \sin \theta \partial \bar{\Omega}^2 / \partial z$ in the middle column. The values are normalized by Ω_0^2 . The last column shows the mean latitudinal entropy gradient $R \nabla_\theta \bar{s} / c_p$. From top to bottom, Run A, Run Ac, and Run B. The white/black dashed line indicate the surface ($r = R$).

radial temperature gradient, which is more important than the radial entropy gradient multiplying the latitudinal temperature gradient.

We also note that, owing to the coronal envelope, differential rotation is able to develop a near-surface-shear layer. In the three simulations we can observe a concentration of contours of $\bar{\Omega}$ near the surface, which indicates the presence of strong shear; see Figure 4. However, only in Runs A, and Ac, the strong shear is below or at the surface, while in Run B, it is concentrated at around $r = 1.1 R$. In our simulation of Runs A and Ac the shear layer is more extended than in the Sun and penetrates deeper into the convection zone. Further




FIG. 6.— Off-diagonal component $\chi_{\theta r}$ and the radial component χ_{rr} of the turbulent heat conductivity tensor normalized by $\chi_{t0} = u_{rms}/3k_f$ and calculated from Equations (22) and (23). From left to right, Run A, Run Ac and Run B. Note, that the high values at the bottom of the convection zone, are due to the vanishing radial entropy gradient. Otherwise the legend is similar to Figure 5.

studies using higher stratification should prove if this is just an artifact of weak stratification, as one might expect. The spoke-like rotation profile with the strong shear near the surface occurs mostly at lower latitudes, i.e. close to the equator. At latitudes above $\pm 30^\circ$ the contours of constant rotation are more complex, but show some indication of strong shear close to the surface and only in Run B the contours become cylindrical beyond $\pm 60^\circ$ latitude. The location of the spoke-like differential rotation profile coincides with a similarly shaped mean latitudinal entropy gradient. The entropy gradient in the northern (southern) hemisphere is negative (positive) below $\pm 30^\circ$ latitude. In Run B, this region reaches to higher latitudes than in Runs A and Ac, which leads to radial contours of angular velocity at higher latitudes.

As mentioned above, we expect that latitudinal entropy gradient to be a consequence of an anisotropic convective diffusivity tensor. Such anisotropies are caused by the rotational influence on the turbulence. In particular, there is a term proportional to $\Omega_{0i}\Omega_{0j}$ (Ω_{0i} is the i th component of Ω_0 , and not to be confused with $\bar{\Omega}(r, \theta)$), which gives a symmetric contribution $\chi_{r\theta} = \chi_{\theta r}$ proportional to $\cos \theta \sin \theta$, so it vanishes at the poles and at the equator. In the presence of a latitudinal entropy gradient, it would lead to an additional contribution to the radial convective flux,

$$\bar{F}_r = -\chi_{rr}\bar{\rho}\bar{T}\nabla_r\bar{s} - \chi_{r\theta}\bar{\rho}\bar{T}\nabla_\theta\bar{s}. \quad (19)$$

Since $\chi_{r\theta} = \chi_{\theta r}$, and since there is a radial entropy gradient,

it would also lead to a contribution in the latitudinal flux,

$$\bar{F}_\theta = -\chi_{\theta r} \bar{\rho} \bar{T} \nabla_r \bar{s} - \chi_{\theta\theta} \bar{\rho} \bar{T} \nabla_\theta \bar{s}. \quad (20)$$

If we ignore the second term proportional to $\nabla_\theta \bar{s}$, we could estimate $\chi_{\theta r}$ by measuring

$$\bar{F}_\theta = c_P \bar{\rho} \overline{u'_\theta T'}, \quad (21)$$

so

$$\chi_{\theta r} \approx -c_P \overline{u'_\theta T'} / \bar{T} \nabla_r \bar{s}. \quad (22)$$

The result is shown in Figure 6, where we also plot a similar estimate of the radial component,

$$\chi_{rr} \approx -c_P \overline{u'_r T'} / \bar{T} \nabla_r \bar{s}. \quad (23)$$

We normalize both by $\chi_{t0} = u_{\text{rms}}/3k_f$ and find that the latter is about 3 in those units, and the former one is about 1.

In reality, and as a consequence, we cannot neglect the second term proportional to $\nabla_\theta \bar{s}$. In fact, as shown in the mean field calculations of Brandenburg et al. (1992), it will try to balance the first term so as to reduce the latitudinal heat flux and thus to produce a latitudinal entropy gradient and thus a baroclinic term as we see it.

Another important result is the solar-like meridional circulation in the convection zone. In the middle column and right column of Figure 4 we plot the meridional circulation in terms of the mass flux as vectors of $\bar{\rho}(\bar{u}_r, \bar{u}_\theta, 0)$ and as radial cuts of $\bar{\rho} \bar{u}_\theta$ through three values of colatitude $\theta_1 = 60^\circ$, $\theta_2 = 75^\circ$, and $\theta_3 = 83^\circ$, corresponding to 30° , 15° , 7° latitudes. Note here that in the northern (southern) hemisphere a negative $\bar{\rho} \bar{u}_\theta$ means poleward (equatorward) flow, and a positive one equatorward (poleward) flow. Runs A and Ac shows significant solar-like meridional circulation, while Run B more diffuse pattern. Looking at Run A, in the northern hemisphere at lower latitudes ($\leq 20^\circ$) just below the surface $r = r_1 = 0.97 R$, the meridional circulation is poleward with $\bar{\rho} \bar{u}_\theta = -0.007 \rho_0 u_{\text{rms}}$. Above the surface there is a return flow in the equatorward direction. This return flow peaks above the surface with a similar flux. The turning point $\bar{\rho} \bar{u}_\theta = 0$ is just below the surface, at around $r = 0.985 R$. If we were to redefine the surface of the simulated star to this radius, we would obtain a solar-like meridional circulation. In the southern hemisphere the pattern is the same, however the meridional circulation is a bit weaker. This behavior can be found also in Run Ac, where the flows are weaker and the turning point lies slightly deeper. Note that the larger return flow is a cause of our particular setup, which has a much lower stratification than the Sun, see Figure 1. Higher stratification should lead to a much weaker return flow above the surface. In Run B, a poleward flow develops in the northern hemisphere only close to equator ($\theta = \theta_3$) or deeper in the convection zone at high latitudes ($\theta = \theta_1$). However, in the southern hemisphere, the pattern is similar to the slower rotating runs, but with weaker circulation. The meridional circulation has a latitudinal dependence. In Runs A and Ac the return flow reaches higher velocities at lower latitudes ($\leq \pm 20^\circ$). The same is valid for the actual poleward circulation below the surface. In Run B, we find the opposite and the return flow increases with latitude, and so does the meridional circulation.

In the right panel of Figure 4, we can also investigate the amount of meridional circulation cells for the lower latitudes. We find that there are at least two cells in the convection zone. In Run A, there is one cell with poleward flow maxima around $r = r_1$ and minima around $r = 0.91 R$, where the mass flux

closer to the surface is as large as in the return flow in the convection zone. A second one is deeper down in the convection zone and has a similar extent and flux. In Run Ac, the pattern is different and the flux is weaker, but there seems to be an indication of a third cell. Note that the stratification leads to stronger mass fluxes over a smaller cross-sectional area deeper in the convection zone than near the surface, while the velocity is similar. In Run B, there are two strong cells of meridional circulation. The cells seem to be more cylindrical than latitudinal, which can also be seen in the phase shift of the pattern for different latitudes. Even deeper down in the convection zone, the meridional flow is much stronger than in Runs A and Ac. This is not surprising: rapidly rotating stars are expected to have a weaker meridional circulation than slower rotators (Köhler 1970).

3.2. Mean magnetic field evolution

The turbulent helical motions generated by convective heat transport together with differential rotation produce a large-scale magnetic field inside the convection zone. It grows exponentially and shows an initial saturation after around $t/\tau = 100$ for Run A, where it is still growing but slowly and linearly. Run B shows a more peculiar behavior; the field seems to have saturated at around $t/\tau = 300$, but it starts growing again at around $t/\tau = 700$ and has not yet saturated. The second growth is possibly related to the change of the oscillatory mode into stable one; see Figure 7 and below. The magnetic and fluid Reynolds numbers of Run B are higher, which should lead to a higher growth rate. However, the rotation rate measured by Co is around half that of Run A, which leads to a slower amplification of the field. At a later time, around $t/\tau = 1000$, the field of Run B becomes comparable to or even stronger than that of Run A. The B_{rms} reaches around 0.5 of the equipartition field strength in Runs A, and Ac, and 0.6 in Run B; see Table 1. The $B_{\text{rms}}^{\text{tot}}$ is around 20% lower, because the field is mainly concentrated in the convection zone. Comparing these results with those of Käpylä et al. (2012) without a corona but otherwise comparable setup, the magnetic field in the current simulations is significantly weaker. Additionally the growth rate of the dynamo is greater than in the models without the corona, where it takes up to five time longer to reach dynamically important field strengths. This is not surprising because the dynamo in the two-layer model is less restricted and has more freedom for different dynamo modes to be excited. There is no restriction due to the magnetic boundary at the surface, which is open in our simulations and restricted to vertical field in the convection zone simulations of Käpylä et al. (2012). This could explain the fast growth in the beginning, but not the decreased saturation level. At the moment the field is weaker, but in particular Run B shows a strong growth starting again at a later time, which is comparable with the time of the runs of Käpylä et al. (2012). This might lead to a new saturation level that is as strong as in the runs of Käpylä et al. (2012). On the other hand, the runs in this work and the runs of Käpylä et al. (2012, 2013) have also differences in other parameters, such as stratification, rotation rate Co or Reynolds numbers Re , Rm , so direct comparison might not be always possible.

Looking just at the B_{rms} in the convection zone or the $B_{\text{rms}}^{\text{tot}}$ in the whole domain in Figure 2, we find no clear evidence of cyclic behavior of the field. However, investigating the different components of the mean magnetic field we find signs of clear oscillatory behavior. In Figure 7, we plot the azimuthal mean magnetic field $\bar{B}_\phi$ and the radial mean field




FIG. 7.— Time evolution of the mean magnetic field in the convection zone. From left to right we show the Runs A, Ac, and B and from top to bottom $\overline{B}_\phi$ and $\overline{B}_r$ at $r = r_1 = 0.97R$ and at $r = r_2 = 0.84R$. Dark blue shades represent negative and light yellow positive values. The dashed horizontal lines show the location of the equator at $\theta = \pi/2$. The magnetic field is normalized by its equipartition value, B_{eq} .




FIG. 8.— Time series of eight snapshots of the mean azimuthal magnetic field $\overline{B}_\phi$ separated by 22 turnover times and covering one full magnetic cycle. Dark blue shades represent negative and light yellow positive values. The dashed line indicates the surface ($r = R$). The field is normalized by the equipartition field strength B_{eq} .

$\overline{B}_r$ over time and latitude at the different heights r_1 and r_2 for Runs A, Ac, and B. The structure of the magnetic field changes as the dynamo evolves from the kinematic regime, where the magnetic field is weak and does not significantly influence the flow. In Run A, the azimuthal and radial mean fields migrate poleward in the kinematic regime close to the equator. The cycle frequency is small, just around $t/\tau = 20$. In Run B, we found a similar behavior. The fast poleward migration happens at lower latitudes ($\pm 40^\circ$) for both runs. We recall that in Run A after a short time ($t/\tau \sim 100$) the field is strong enough to backreact on the flow. Now two things happens simultaneously: an oscillating mean magnetic field is starting migrating equatorward at higher latitudes and the fast poleward migration becomes slower. The period of the equa-

torward oscillation is longer and is between $t/\tau = 100$ and 150. The poleward migration near the equator slows down until it finally turns into an equatorward migration aligned with the migration at the higher latitudes ($t/\tau = 500$). Thus we have equatorward migration of the mean radial and azimuthal fields at all latitudes until around $t/\tau = 1500$. At this time the dynamo mode changes, and consequently its latitudinal migration pattern changes. The equatorward migrating and oscillating field near the poles stays stable during whole simulation, but near the equator the field changes. In the northern hemisphere there is a transient poleward migration, which is in phase with the equatorward migration near the poles. In the southern hemisphere the equatorward migration is still dominant, but a stationary mode is superposed near the equator.

The migration patterns are not just features appearing close to the surface, but they penetrate the whole convection zone until the bottom, as seen in the two bottom rows of Figure 7. Therefore, we can rule out that the meridional circulation is the main driver of this migration. As discussed in Section 3.1, the meridional circulation shows strong variability in radius and has at least two cells.

Run Ac has been started from a snapshot of Run A after around $t/\tau = 1150$ by increasing the magnetic Prandtl number; see Table 1. The pattern of the mean magnetic field is not strongly effected by this change of parameter and is similar to Run A. There is a clear equatorward migration near the poles and some indication of poleward migration and equatorward migration near the equator in the northern hemisphere. The stationary fields superimposed to this cycle are similar to those in Run A.

In Run B, where the fluid and magnetic Reynolds numbers are higher and the rotation slower than in Run A, the structure of the mean field evolution shows some differences. In the kinematic regime, the field is similar to that of Run A in which it migrates poleward at lower latitudes. Also, as the field gets stronger it begins to migrate from higher latitudes toward the equator and the fast poleward branch becomes slower. The main difference to Run A is that the poleward migration does not turn into equatorward migration near the equator. In Runs A and Run Ac, the field strength has no clear latitudinal dependence. But in Run B the field strength near the poles is around the half the strength than near the equator. Only during late times the polar branch increases in strength. In Runs A and Ac, the radial and azimuthal components have approximately the same strength, whereas in Run B, the radial mean magnetic field seems to be weaker by a factor of two. Also, Run B shows no clear radial dependence in the structure of the mean field. The period of the equatorward migrating field is around $t/\tau = 200$, which is a bit longer than for Run A. The poleward migration near the equator has an irregular oscillation and is usually not in phase with the equatorward migration near the poles. At around $t/\tau = 1000$ the dynamo mode changes significantly. Not only the magnetic field starts to grow (see Figure 2), but also the magnetic field changes from an oscillatory pattern to a stationary one, or at least an oscillatory one with a much longer period; see Figure 7. In particular in the norther hemisphere the mean azimuthal field shows a strong increase in strength. The field pattern now consists of a strong time-independent field with a latitudinal dependence. Near the surface (at $r = r_1$; see Section 2) the field seems to migrate slowly toward the equator, but it is not possible to identify a migrating pattern in the present run.

If one translates the cycle period of the equatorward migration to solar values using of turnover time τ of 1 month, we obtain a cycle period of 12 years and 16 years for Runs A and B, respectively. This would be a typical value in the middle of the convection zone.

To investigate the equatorward migration in the first half of the simulations, we plot the mean azimuthal magnetic field $\overline{B}_\phi$ for eight different times for Run A, resolving one cycle; see Figure 8. The field penetrates the whole convection zone and has up to four regions with different polarities in one hemisphere. These polarities are migrating toward the equator. In the northern hemisphere at around 45° latitude, there is a magnetic field concentration with positive polarity. After $\Delta t/\tau = 87$ (5th panel), at the same location we find a nega-

tive magnetic field concentration, and again after $\Delta t/\tau = 65$ (8th panel) the same polarity appear as in the beginning of the cycle. One can see a clear cyclic equatorward migration of the field, but it is irregular. The two hemispheres do not show the same magnetic field strength and it seems that, from time to time, there is only one dominant polarity in one hemisphere, while in the other there are three. Note also the strong negative magnetic fields near and above the surface, which seem to show also an cycle dependency.

4. CONCLUSIONS

We have used a model that combines the turbulent convective dynamo with a coronal layer to reproduce properties of the Sun. We found a solar-like differential rotation with roughly radial contours of angular velocity at low latitudes. This is accompanied by poleward meridional circulation just below the surface and a return flow above the surface. Additionally the differential rotation profiles show a near-surface-shear layer in two of the three simulations we perform. In the third simulation, this layer is still exists but it is now located above the surface. We identify as the main driver of spoke-like differential rotation the non-zero latitudinal entropy gradient, which is self-consistently generated.

The mean magnetic field shows a pattern of equatorward migration similar to Käpylä et al. (2012, 2013). This pattern is mostly visible at higher latitudes and only in two of the simulation to lower latitudes. However, in later stages of the simulation the equatorward migration is only visible at high latitudes and at lower latitudes poleward migration or stationary modes occur. In one of the simulation the dynamo mode changed completely to stationary mode on all latitudes, at later stages. The dynamo has a shorter excitation time than in the earlier work of Käpylä et al. (2012).

This work leads to the conclusion that the inclusion of a coronal layer to a convective dynamo simulation has an influence on the fluid and magnetic properties of the interior. In recent simulations, we were able to produce recurrent coronal ejections from the solar surface (Warnecke et al. 2012b) using a two layer approach. In earlier models of forced turbulence with a coronal layer (Warnecke & Brandenburg 2010; Warnecke et al. 2011, 2012a), we also found the ejection of magnetic helicity out of the dynamo region. These ejections can support and amplify the magnetic field due to enhanced helicity fluxes.

Here, we present evidence that even the fluid properties in the bulk of the convection zone might be influenced by the coronal layer. Solar-like rotation profiles could not be reproduced by earlier direct numerical simulation of convective dynamo (Käpylä et al. 2012) without prescribing a latitudinal entropy gradient at the bottom of the convection zone (Miesch et al. 2006). However, to have more convincing evidence in support of this, we need to perform a detailed parameter study using different coronal sizes and compare them with simulations without a corona.

Another extension of our work would be the measurement of magnetic helicity fluxes through the surface and how their dependence on the coronal size. To investigate the mechanism of the equatorward migration, which is crucial for understanding the solar dynamo, one should measure the turbulent transport coefficients through approaches like the test-field method (Schrinner et al. 2007).

In further work we will investigate the possibility of producing coronal ejections using the setup of these runs. In comparison with Warnecke et al. (2012b), we use here a corona

with a much higher temperature, and a lower plasma beta. It will be interesting to see how the coronal ejections are influenced by this.

We acknowledge the allocation of computing resources provided by the Swedish National Allocations Committee at the Center for Parallel Computers at the Royal Institute of Technology in Stockholm, the National Supercomputer Centers in Linköping and the High Performance Computing Center North in Umeå. Part of the computations have been carried out in the facilities hosted by the CSC – IT Center for Sci-

ence in Espoo, Finland, which are financed by the Finnish ministry of education. This work was supported in part by the European Research Council under the AstroDyn Research Project No. 227952, the Swedish Research Council Grant No. 621-2007-4064, and the Academy of Finland grants 136189, 140970 (PJK) and 218159, 141017 (MJM), University of Helsinki ‘Active Suns’ research project, as well as the HPC-Europa2 project, funded by the European Commission - DG Research in the Seventh Framework Programme under grant agreement No. 228398. The authors thank NORDITA for hospitality during their visits.

REFERENCES

- Blackman, E. G., & Brandenburg, A. 2003, *ApJ*, 584, L99
 Brandenburg, A. 2005, *ApJ*, 625, 539
 Brandenburg, A., Moss, D., & Tuominen, I. 1992, *A&A*, 265, 328
 Brandenburg, A., & Sandin, C. 2004, *A&A*, 427, 13
 Brandenburg, A., & Subramanian, K. 2005, *Phys. Rep.*, 417, 1
 Brown, B. P., Miesch, M. S., Browning, M. K., Brun, A. S., & Toomre, J. 2011, *ApJ*, 731, 69
 Brun, A. S., Miesch, M. S., & Toomre, J. 2004, *ApJ*, 614, 1073
 Dikpati, M., & Charbonneau, P. 1999, *ApJ*, 518, 508
 Ghizaru, M., Charbonneau, P., & Smolarkiewicz, P. K. 2010, *ApJ*, 715, L133
 Gilman, P. A. 1983, *ApJS*, 53, 243
 Hubbard, A., & Brandenburg, A. 2012, *ApJ*, 748, 51
 Käpylä, P. J., Korpi, M. J., Brandenburg, A., Mitra, D., & Tavakol, R. 2010, *Astron. Nachr.*, 331, 73
 Käpylä, P. J., Korpi, M. J., & Tuominen, I. 2006, *Astron. Nachr.*, 327, 884
 Käpylä, P. J., Mantere, M. J., & Brandenburg, A. 2011a, *Astron. Nachr.*, 332, 883
 —. 2012, *ApJ*, 755, L22
 Käpylä, P. J., Mantere, M. J., Cole, E., Warnecke, J., & Brandenburg, A. 2013, *ApJ*, submitted, arXiv and Nordita preprint 2013-4
 Käpylä, P. J., Mantere, M. J., Guerrero, G., Brandenburg, A., & Chatterjee, P. 2011b, *A&A*, 531, A162
 Karak, B. B. 2010, *ApJ*, 724, 1021
 Kitchatinov, L. L., & Olemskoy, S. V. 2012, *Sol. Phys.*, 276, 3
 Kitchatinov, L. L., & Rüdiger, G. 1995, *A&A*, 299, 446
 Köhler, H. 1970, *Sol. Phys.*, 13, 3
 Krause, F., & Rädler, K.-H. 1980, *Mean-field Magnetohydrodynamics and Dynamo Theory* (Oxford: Pergamon Press)
 Miesch, M. S., Brun, A. S., & Toomre, J. 2006, *ApJ*, 641, 618
 Mitra, D., Tavakol, R., Brandenburg, A., & Moss, D. 2009, *ApJ*, 697, 923
 Rüdiger, G. 1980, Reynolds stresses and differential rotation. I - On recent calculations of zonal fluxes in slowly rotating stars, Vol. 16, 239–261
 —. 1989, *Differential Rotation and Stellar Convection. Sun and Solar-type Stars* (Berlin: Akademie Verlag)
 Schrunner, M., Rädler, K.-H., Schmitt, D., Rheinhardt, M., & Christensen, U. R. 2007, *Geophys. Astrophys. Fluid Dyn.*, 101, 81
 Spiegel, E. A., & Weiss, N. O. 1980, *Nature*, 287, 616
 Warnecke, J., & Brandenburg, A. 2010, *A&A*, 523, A19
 Warnecke, J., Brandenburg, A., & Mitra, D. 2011, *A&A*, 534, A11
 —. 2012a, *J. Spa. Weather Spa. Clim.*, 2, A11
 Warnecke, J., Käpylä, P. J., Mantere, M. J., & Brandenburg, A. 2012b, *Sol. Phys.*, 280, 299